arXiv:math/0107161v1 [math.AG] 23 Jul 2001

Simpson Jacobians of generalized tree-like curves

Ana Cristina López Martín

February 1, 2008

Abstract

The compactified Jacobian of any projective curve X is defined as the Simpson moduli space of torsion free rank one degree d sheaves that are semistable with respect to a fixed polarization H on X. In this paper we give explicitly the structure of this compactified Simpson Jacobian in the case where X is a generalized tree-like curve, i.e., a projective, reduced and connected curve such that the intersection points of its irreducible components are disconnecting ordinary double points. We prove that it is isomorphic to the product of the compactified Jacobians of a certain degree d_i of its components C_i, where the degrees d_i depend on d, H and on the particular structure of the curve. We find also necessary and sufficient conditions for the existence of stable points which allow us to study the variation of these Simpson Jacobians as the polarization H changes.

1 Introduction

The problem of compactifying the generalized Jacobian of a singular curve has been studied since Igusa's work [I] around 1950. He constructed a compactification of the Jacobian of a nodal and irreducible curve X as the limit of the Jacobians of smooth curves approaching X. Igusa also showed that his compactification does not depend on the considered family of smooth curves. An intrinsic characterization of the boundary points of the Igusa's compactification as the torsion free, rank 1 sheaves which are not line bundles is due to Mumford and Mayer. The complete construction for a family of integral curves over a noetherian Hensel local ring with residue field separably closed was carried out by D'Souza [D'So]. One year later, Altman and Kleiman [AK] gave the construction for a general family of integral curves.

When the curve X is reducible and nodal, Oda and Seshadri [OS] produced a family of compactified Jacobians Jac_ϕ parameterized by an element ϕ of a real vector space. Seshadri dealt in [Ses] with the general case of a reduced curve considering sheaves of higher rank as well.

In 1994, Caporaso showed [C] how to compactify the relative Jacobian over the moduli of stable curves and described the boundary points of the compactified Jacobian of a stable curve X as invertible sheaves on certain Deligne-Mumford semistable curves that have X as a stable model. Recently, Pandharipande [P] has given another construction with the boundary points now representing torsion free, rank 1 sheaves and he showed that the Caporaso's compactification was equivalent to his.

On the other hand, Esteves [E] constructed a compactification of the relative Jacobian of a family of geometrically reduced and connected curves and compared it with Seshadri's construction [Ses] using theta funtions and Alexeev [A] gave a description of the Jacobian of certain singular curves in terms of the orientations on complete subgraphs of the dual graph of the curve.

Simpson's work [S] on the moduli of pure coherent sheaves on projective spaces allows us to define in a natural way the Jacobian of any projective curve X as the space $\mathrm{Jac}^d(X)_s$ of equivalence classes of stable invertible sheaves with degree d. This is precisely the definition we adopt and we also denote by $\overline{\mathrm{Jac}}^d(X)$ the space of equivalence classes of semistable, pure dimension 1, rank 1 sheaves with degree d.

In some recent papers about the moduli spaces of stable vector bundles on elliptic fibrations, for instance [HM], the Jacobian in the sense of Simpson of spectral curves appears. Beauville [B] uses it as well in counting the number of rational curves on K3 surfaces. This suggests the necessity to determine the structure of these Simpson Jacobians.

The aim of this work is to describe the Simpson Jacobian of a generalized tree-like curve, that is, a projective, reduced and connected curve such that the intersection points of its irreducible components are disconnecting ordinary double points.

Let H be the fixed polarization on X. Let $C_1, \dots, C_N$ denote the irreducible components of X and let $P_1, \dots, P_{N-1}$ be the intersection points of

$C_1, \ldots, C_N$. The first result we use is a lemma of Teixidor [T] that allows to order C_i and to find subcurves X_i of X that are also generalized tree-like curves and intersect its complement in X at just one point P_i. Then, we write $k_{X_i} = \frac{h_{X_i}(b+1)}{h}$ for $i = 1, \ldots, N-1$, where h is the degree of H, h_{X_i} is the degree of the induced polarization on X_i and b is the residue class of d minus the arithmetic genus of X modulo h.

The theorem that gives the description of the schemes $\operatorname{Jac}^d(X)_s$ and $\overline{\operatorname{Jac}}^d(X)$ is the following:

Theorem 3.1. *Let* $X = C_1 \cup \ldots \cup C_N$, $N \geq 2$, *be a generalized tree-like curve with a polarization* H. *If* k_{X_i} *is not integer for* $i \leq N-1$, *then* $\operatorname{Jac}^d(X)_s$ *is equal to* $\prod_{i=1}^N \operatorname{Pic}^{d_i^X}(C_i)$ *and*

$$\operatorname{Jac}^d(X)_s \subseteq \overline{\operatorname{Jac}}^d(X)_s = \overline{\operatorname{Jac}}^d(X) \simeq \prod_{i=1}^N \overline{\operatorname{Jac}}^{d_i^X}(C_i)$$

where d_i^X *are integer numbers inductively constructed.*

If k_{X_i} *is integer for some* $i \leq N-1$, *then* $\operatorname{Jac}^d(X)_s$ *and* $\overline{\operatorname{Jac}}^d(X)_s$ *are empty and*

$$\overline{\operatorname{Jac}}^d(X) \simeq \prod_{i=1}^N \overline{\operatorname{Jac}}^{d_i}(C_i)$$

where d_i *are integer numbers recursively constructed with an algorithm.*

As we will see, these integer numbers, d_i and d_i^X, depend only on the degree of H and on the fixed ordering of the irreducible components of X.

The first we prove is that a pure dimension 1 sheaf of rank 1 and degree d on X that is locally free at P_i, $i = 1, \ldots, N-1$, is stable if and only if $k_{X_i} \notin \mathbb{Z}$ for $i = 1, \ldots, N-1$ and F is obtained by gluing torsion free rank 1 sheaves on C_i whose degrees are precisely the integers d_i^X (see lemmas 3.1 and 3.2 if F is a line bundle, otherwise the proof is the same). By the contrary, if F is not locally free at some intersection point P_i, the sheaf cannot be stable. Finally, we prove that strictly semistable torsion free rank 1 sheaves exist only when $k_{X_i} \in \mathbb{Z}$ for some $i = 1, \ldots, N-1$ and in this case we construct a Jordan-Hölder filtration for it to conclude that its S-equivalence class belongs to $\prod_{i=1}^N \overline{\operatorname{Jac}}^{d_i}(C_i)$.

It is important to note that, although $\overline{\operatorname{Jac}}^d(C_i) \simeq \overline{\operatorname{Jac}}^{d'}(C_i)$ for any integers d and d', because C_i is an integral curve, to obtain a point in $\overline{\operatorname{Jac}}^d(X)$ it is necessary to consider sheaves on C_i of degrees exactly the integer numbers d_i

Theorem 3.1 allows also to describe a wall-crossing structure for these Simpson Jacobians, in other words, to analyze the variation of the moduli spaces $\mathrm{Jac}^d(X)_s$ as the polarization changes. The variation of the polarization is studied by considering the variation of the numbers k_{X_i}, $(i = 1, \ldots, N-1)$. The space of possible values of such numbers contains a finite number of walls, the hyperplanes $k_{X_i} = a$ where a is an integer number $1 \leq a \leq h_{X_i}$. The complement of the walls has a finite number of connected components each of them called a chamber. Therefore, Theorem 3.1 means that in the interior of a given chamber the Jacobian $\mathrm{Jac}^d(X)_s$ is not empty and independent of the choice of numbers k_{X_i}. When the numbers k_{X_i} lie in a wall, the theorem shows that the Simpson Jacobian of X becomes empty. Finally, if we choose two families of numbers k_{X_i} lying in the interior of two different chambers, then the structure of the corresponding Simpson Jacobians is the same but the degrees d_i^X of the induced line bundles on every component C_i change according to (1).

If α is a real number, we use $[\alpha]$ to denote the greatest integer less than or equal to α.

2 Preliminaries

Let X be a projective, reduced and connected curve over an algebraically closed field κ. Let $C_1, \ldots, C_N$ denote the irreducible components of X. Let $\mathcal{L}$ be an ample invertible sheaf on X, let H be the associated polarization and let h be the degree of H.

Let F be a coherent sheaf on X. We say that F is pure of dimension one or torsion free if for all nonzero subsheaves $F' \hookrightarrow F$ the dimension of $\mathrm{Supp}(F')$ is 1. The rank and the degree (with respect to H) of F are the rational numbers $\mathrm{r}_H(F)$ and $\mathrm{d}_H(F)$ determined by the Hilbert polynomial

$$P(F, n, H) = \chi(F \otimes \mathcal{O}_X(nH)) = h\,\mathrm{r}_H(F)n + \mathrm{d}_H(F) + \mathrm{r}_H(F)\chi(\mathcal{O}_X).$$

The slope of F is defined by

$$\mu_H(F) = \frac{\mathrm{d}_H(F)}{\mathrm{r}_H(F)}$$

The sheaf F is stable (resp. semistable) with respect to H if F is pure of dimension one and for any proper subsheaf $F' \hookrightarrow F$ one has

$$\mu_H(F') < \mu_H(F) \text{ (resp. } \leq)$$

In [S] Simpson defined the rank as the integer number $h\,\mathrm{r}_H(F)$ and the slope as the quotient

$$\frac{\mathrm{d}_H(F) + \mathrm{r}_H(F)\chi(\mathcal{O}_X)}{h\,\mathrm{r}_H(F)}$$

Stability and semistability considered in terms of Simpson's slope and in terms of μ_H are equivalent. We adopt these definitions of rank and degree of F because they coincide with the classical ones when the curve is integral.
Notation: We shall use the following notation. For every proper subcurve D of X, we will denote by F_D the restriction of F to D modulo torsion, that is, $F_D = (F \otimes \mathcal{O}_D)/\text{torsion}$, π_D will be the surjective morphism $F \to F_D$ and $F^D = \ker \pi_D$. We shall denote by h_D the degree of the induced polarization H_D on D. If $d = \mathrm{d}_H(F)$ then we shall write $d_D = \mathrm{d}_{H_D}(F_D)$.

We first recall some general properties we will use later.

Lemma 2.1. *Let F be a pure dimension one sheaf on X supported on a subcurve D of X. Then F is stable (resp. semistable) with respect to H_D if and only if F is stable (resp. semistable) with respect to H.*

Proof. It follows from the equality

$$P(F,n,H) = \chi(i_*F \otimes \mathcal{O}_X(nH)) = \chi(F \otimes \mathcal{O}_D(nH_D)) = P(F,n,H_D)$$

where $i\colon D \hookrightarrow X$ is the inclusion map. $\square$

Lemma 2.2. *A torsion free rank 1 sheaf F on X is stable (resp. semistable) if and only if $\mu_H(F^D) < \mu_H(F)$ (resp. $\leq$) for every proper subcurve D of X.*

Proof. Given a subsheaf G of F such that $\mathrm{Supp}(G) = D \subset X$, let us consider the complementary subcurve $\overline{D}$ of D in X, i.e. the closure of $X - D$. Since $F_{\overline{D}}$ is torsion free, we have $G \subset F^{\overline{D}}$ with $\mathrm{r}_H(G) = \mathrm{r}_H(F^{\overline{D}})$ so that $\mu_H(G) \leq \mu_H(F^{\overline{D}})$ and the result follows. $\square$

Let $P_1, \dots, P_k$ denote the intersection points of $C_1, \dots, C_N$. It is known (see [Ses]) that for every pure dimension one sheaf F on X there is an exact sequence

$$0 \to F \to F_{C_1} \oplus \ldots \oplus F_{C_N} \to T \to 0$$

where T is a torsion sheaf whose support is precisely the set of those points P_i, $i = 1, \dots, k$, where F is locally free. From this exact sequence, we get

$$\begin{aligned}
\mathrm{r}_H(F) &= \frac{1}{h} \sum_{i=1}^{N} h_{C_i} \, \mathrm{r}_{H_{C_i}}(F_{C_i}) \\
\mathrm{d}_H(F) &= \sum_{i=1}^{N} (\mathrm{d}_{H_{C_i}}(F_{C_i}) + \mathrm{r}_{H_{C_i}}(F_{C_i}) \chi(\mathcal{O}_{C_i})) - \mathrm{r}_H(F) \chi(\mathcal{O}_X) - \chi(T).
\end{aligned}$$

In particular, if F is a torsion free sheaf of rank 1 with respect to H, then for every proper subcurve D of X F_D is torsion free of rank 1 with respect to H_D.

The following lemma, also due to Seshadri, describes the stalk of a torsion free sheaf on X at the points P_i.

Lemma 2.3. *Let F be a pure dimension one sheaf on X. If P_i is an ordinary double point lying in two irreducible components C_i^1 and C_i^2, then*

$$F_{P_i} \simeq \mathcal{O}_{X,P_i}^{a_1} \oplus \mathcal{O}_{C_i^1,P_i}^{a_2} \oplus \mathcal{O}_{C_i^2,P_i}^{a_3}$$

where $a_1,\ a_2,\ a_3$ are the integer numbers determined by:

$$\begin{aligned}
a_1 + a_2 &= \mathrm{rk}(F_{P_i} \underset{\mathcal{O}_{X,P_i}}{\otimes} \mathcal{O}_{C_i^1,P_i}) \\
a_1 + a_3 &= \mathrm{rk}(F_{P_i} \underset{\mathcal{O}_{X,P_i}}{\otimes} \mathcal{O}_{C_i^2,P_i}) \\
a_1 + a_2 + a_3 &= \mathrm{rk}(F_{P_i} \otimes \kappa)
\end{aligned}$$

Proof. See [Ses], Huitiême Partie, Prop. 3. ☐

According to the general theory, for every semistable sheaf F with respect to H there is a Jordan-Hölder filtration

$$0 = F_0 \subset F_1 \subset \ldots \subset F_n = F$$

with stable quotients F_i/F_{i-1} and $\mu_H(F_i/F_{i-1}) = \mu_H(F)$ for $i = 1, \dots, n$. This filtration need not be unique, but the graded object $Gr(F) = \bigoplus_i F_i/F_{i-1}$

does not depend on the choice of the Jordan-Hölder filtration. Two semistable sheaves F and F' on X are said to be S-equivalent if $Gr(F) \simeq Gr(F')$. Note that two stable sheaves are S-equivalent only if they are isomorphic.

In the relative case, given a scheme S of finite type over κ, a projective morphism of schemes $f\colon X \to S$ whose geometric fibers are curves and a relative polarization H, we define the relative rank and degree of a coherent sheaf F on X, flat over S, as its rank and degree on fibers, and we say that F is relatively pure of dimension one (resp. stable, resp. semistable) if it is flat over S and if its restriction to every geometric fiber of f is pure of dimension one (resp. stable, resp. semistable).

Let $\mathbf{Jac^d(X/S)_s}$ (resp. $\overline{\mathbf{Jac}}^{\mathbf{d}}\mathbf{(X/S)_s}$) be the functor which to any S-scheme T associates the set of equivalence classes of stable invertible (resp. relatively torsion free rank 1) sheaves on $X_T = X \underset{S}{\times} T$ with relative degree d. Two such sheaves F and F' are said to be equivalent if $F' \simeq F \otimes f_T^* N$, where N is a line bundle on T and $f_T\colon X_T \to T$ is the natural projection. Similarly, we define the functor $\mathbf{Jac^d(X/S)}$ (resp. $\overline{\mathbf{Jac}}^{\mathbf{d}}\mathbf{(X/S)}$) of semistable invertible (resp. relatively torsion free rank 1) sheaves.

As a particular case of the Simpson's work [S], there exists a projective scheme $\overline{\operatorname{Jac}}^d(X/S) \to S$ which coarsely represents the functor $\overline{\mathbf{Jac}}^{\mathbf{d}}\mathbf{(X/S)}$. Rational points of $\overline{\operatorname{Jac}}^d(X/S)$ correspond to S-equivalence classes of semistable torsion free sheaves of rank 1 and degree d on a fiber X_s ($s \in S$). Moreover, $\mathbf{Jac^d(X/S)}$ is coarsely represented by a subscheme $\operatorname{Jac}^d(X/S)$ and there are open subschemes $\operatorname{Jac}^d(X/S)_s$ and $\overline{\operatorname{Jac}}^d(X/S)_s$ which represent the other two funtors.

Definition 2.1. *The Simpson Jacobian of* X *is* $\operatorname{Jac}^d(X)_s = \operatorname{Jac}^d(X/\operatorname{Spec}\kappa)_s$. *We denote* $\overline{\operatorname{Jac}}^d(X) = \overline{\operatorname{Jac}}^d(X/\operatorname{Spec}\kappa)$ *and* $\overline{\operatorname{Jac}}^d(X)_s = \overline{\operatorname{Jac}}^d(X/\operatorname{Spec}\kappa)_s$.

When X is an integral curve every torsion free rank 1 sheaf is stable, and then $\operatorname{Jac}^d(X) = \operatorname{Jac}^d(X)_s$ is equal to the Picard scheme $\operatorname{Pic}^d(X)$ and $\overline{\operatorname{Jac}}^d(X) = \overline{\operatorname{Jac}}^d(X)_s$ coincides with the Altman-Kleiman's compactification [AK].

Definition 2.2. *A generalized tree-like curve is a projective, reduced and connected curve* $X = C_1 \cup \ldots \cup C_N$ *over* κ *such that the intersection points,* $P_1, \ldots, P_k$*, of its irreducible components are disconnecting ordinary double points.*

Henceforth we shall assume that X is a generalized tree-like curve, and then $k = N - 1$.

Lemma 2.4. *It is possible to order the irreducible components* $C_1, \dots, C_N$ *of the curve* X*, so that for every* $i \leq N-1$ *all but one of the connected components of* $X - C_i$ *consists entirely of irreducible components with subindices smaller than* i*. Then there are irreducible components, say* $C_{i_1}, \dots, C_{i_k}$*, with all subindices smaller than* i*, such that* $X_i = C_i \cup C_{i_1} \cup \ldots \cup C_{i_k}$ *is connected and intersects its complement* $\overline{X}_i$ *in* X *in just one point* P_i*.*

Proof. Teixidor proves this lemma in ([T], Lem. 1) when X is a tree-like curve, that is, its irreducible components are smooth, but her proof is valid for our curve. □

3 The Description

Let us suppose from now on that an ordering of the components of X as in lemma 2.4, has been fixed.

Notation: If $g = \mathrm{g}(X)$ denote the arithmetic genus of X, that is, the dimension of $\mathrm{H}^1(X, \mathcal{O}_X)$, for any torsion free rank 1 sheaf F en X of degree d, let b be the residue class of $d - g$ modulo h so that

$$d - g = ht + b.$$

For every proper subcurve D of X, we shall write

$$k_D = \frac{h_D(b+1)}{h}.$$

Lemma 2.4 allows us to define inductively integer numbers d_i^X as follows:

$$\begin{aligned} d_i^X &= -\chi(\mathcal{O}_{X_i}) + h_{X_i} t + [k_{X_i}] + 1 - d_{i_1}^X - \ldots - d_{i_k}^X, \quad \text{for } i = 1, \dots, N-1 \\ d_N^X &= d - d_1^X - \ldots - d_{N-1}^X. \end{aligned} \tag{1}$$

We are now going to modify the above numbers to obtain new numbers d_i associated with X. This is accomplished by a recurrent algorithm. In order to describe it we start by saying that a connected subcurve $D = C_{j_1} \cup \ldots \cup C_{j_r}$, $r \geq 1$, of X ordered according to lemma 2.4 is *final* either when the numbers $k_{D_{j_t}}$ are not integers for $t = 1, \dots, r-1$ or D is irreducible.

If D is a final curve, we define d_{j_t} as follows:

1. if $r > 1$, $d_{j_t} = d^D_{j_t}$ for $t = 1, \ldots, r$.

2. if $r = 1$, $d_{j_1} = h_{C_{j_1}} t + [k_{C_{j_1}}] - \chi(\mathcal{O}_{C_{j_1}})$.

Algorithm: If the curve X is final, $d_i = d^X_i$ for all i. Otherwise, let i be the first index for which $k_{X_i} \in \mathbb{Z}$. We consider the two connected components, $Y = X_i$ and $\overline{Y} = \overline{X_i}$, of $X - P_i$ and we reorder them according with lemma 2.4. This induces a new ordering P^Y_r (resp. $P^{\overline{Y}}_s$) of the points P_j ($j \neq i$) in Y (resp. $\overline{Y}$). Then,

a) If Y (resp. $\overline{Y}$) is a final curve, the process finishes for Y (resp. $\overline{Y}$).

b) If Y is not final, we take the first index r of Y for which $k_{Y_r} \in \mathbb{Z}$, consider the connected components, Z and $\overline{Z}$, of $Y - P^Y_r$ and reorder them according with lemma 2.4. If Z and $\overline{Z}$ are final, the process finishes for Y. Otherwise, we iterate the above argument for those components that are not final and so on. The process finishes for Y when all subcurves that we find are final.

c) If $\overline{Y}$ is not final, we take the first index s of $\overline{Y}$ such that $k_{\overline{Y}_s} \in \mathbb{Z}$, consider the connected components, W and $\overline{W}$, of $\overline{Y} - P^{\overline{Y}}_s$ and reorder them according with lemma 2.4. If W and $\overline{W}$ are final, the process finishes for $\overline{Y}$. Otherwise, we repeat the above argument for those components that are not final and so on. The process finishes for $\overline{Y}$ when all subcurves that we obtain are final.

The algorithm for X finishes when it finishes for both Y and $\overline{Y}$. □

We can now state the theorem that determines the structure of the Simpson Jacobian of X and of the schemes $\overline{\operatorname{Jac}}^d(X)_s$ and $\overline{\operatorname{Jac}}^d(X)$.

Theorem 3.1. *Let $X = C_1 \cup \ldots \cup C_N$, $N \geq 2$, be a generalized tree-like curve.*

a) If k_{X_i} is not an integer for every $i \leq N-1$, then

$$\operatorname{Jac}^d(X)_s = \prod_{i=1}^N \operatorname{Pic}^{d^X_i}(C_i) \quad \textit{and}$$

$$\operatorname{Jac}^d(X)_s \subseteq \overline{\operatorname{Jac}}^d(X)_s = \overline{\operatorname{Jac}}^d(X) \simeq \prod_{i=1}^N \overline{\operatorname{Jac}}^{d^X_i}(C_i)$$

where d^X_i are the above integers.

b) If k_{X_i} is an integer for some $i \leq N-1$, then

$$\operatorname{Jac}^d(X)_s = \overline{\operatorname{Jac}}^d(X)_s = \emptyset \quad \textit{and}$$

$$\overline{\mathrm{Jac}}^d(X) \simeq \prod_{i=1}^N \overline{\mathrm{Jac}}^{d_i}(C_i)$$

where d_i are the integers constructed with the above algorithm.

In order to prove that the Simpson Jacobian of X is not empty only when $k_{X_i} \notin \mathbb{Z}$ for all $i \leq N-1$, we need the next two lemmas that characterize stable invertible sheaves on X.

Lemma 3.1. *Let L be a line bundle on X of degree d. If L is stable, then k_{X_i} is not an integer for every $i \leq N-1$ and L is obtained by gluing invertible sheaves L_i on C_i of degrees d_i^X, $i = 1, \dots, N$.*

Proof. Let us consider the subsheaves L^{X_i} of L, $i = 1, \dots, N-1$, X_i being the subcurves of X given by lemma 2.4. By the stability of L, we get

$$h_{X_i} d - h d_{X_i} \underset{(\leq)}{<} h\chi(\mathcal{O}_{X_i}) - h_{X_i}\chi(\mathcal{O}_X) \quad \text{for } i = 1, \dots, N-1 \tag{2}$$

Considering the subsheaves $L^{\overline{X}_i}$ of L, $i = 1, \dots, N-1$, yields

$$h_{\overline{X}_i} d - h d_{\overline{X}_i} \underset{(\leq)}{<} h\chi(\mathcal{O}_{\overline{X}_i}) - h_{\overline{X}_i}\chi(\mathcal{O}_X) \quad \text{for } i = 1, \dots, N-1 \tag{3}$$

Since $X = X_i \cup \overline{X}_i$ and X_i, $\overline{X}_i$ meet only at P_i, we have $d = d_{X_i} + d_{\overline{X}_i}$, $h = h_{X_i} + h_{\overline{X}_i}$ and $\chi(\mathcal{O}_X) = \chi(\mathcal{O}_{X_i}) + \chi(\mathcal{O}_{\overline{X}_i}) - 1$. Then, (2) and (3) give

$$h_{X_i} d + h_{X_i}\chi(\mathcal{O}_X) - h\chi(\mathcal{O}_{X_i}) \underset{(\leq)}{<} h d_{X_i} \underset{(\leq)}{<} h_{X_i} d + h_{X_i}\chi(\mathcal{O}_X) - h\chi(\mathcal{O}_{X_i}) + h \tag{4}$$

We have $\mathrm{r}_{H_{X_i}}(L_{X_i}) = 1$ so that $d_{X_i} = \mathrm{d}_{H_{X_i}}(L_{X_i})$ is an integer. Then, if $k_{X_i} \in \mathbb{Z}$ for some $i \leq N-1$, (4) becomes a contradiction. Thus $k_{X_i} \notin \mathbb{Z}$ for all $i \leq N-1$ and there is only one possibility for d_{X_i}, namely

$$d_{X_i} = -\chi(\mathcal{O}_{X_i}) + h_{X_i} t + [k_{X_i}] + 1, \qquad \text{for } i = 1, \dots, N-1.$$

From $d_{X_i} = d_{C_i} + d_{C_{i_1}} + \ldots + d_{C_{i_k}}$ and the exact sequence

$$0 \to L \to L_{C_1} \oplus \ldots \oplus L_{C_N} \to \oplus_{i=1}^{N-1} \kappa(P_i) \to 0$$

we deduce that $d_{C_i} = d_i^X$ for all i and the proof is complete. □

The following lemma, which is similar to lemma 2 of Teixidor [T], proves the converse of lemma 3.1.

Lemma 3.2. *Let L be an invertible sheaf of degree $d = g + ht + b$ obtained by gluing line bundles L_i on C_i of degrees d_i^X, $i = 1, \ldots, N$. Suppose that k_{X_i} is not integer for every $i \leq N - 1$ and let $D = C_{a(1)} \cup \ldots \cup C_{a(t)}$ be a proper subcurve of X. Then,*

1. The following inequality holds:

$$-\chi(\mathcal{O}_D) + h_D t + k_D < d_D < -\chi(\mathcal{O}_D) + h_D t + k_D + \alpha$$

where α is the number of intersection points of D and its complement $\overline{D}$ in X.

2. L is stable.

Proof. 1. If D is a connected subcurve of X, let us denote by $C_{i(k)}$, $k = 1, \ldots, \alpha$, (resp. $C_{j(k)}$) the component of D (resp. $\overline{D}$) which contains the point P_k (it is possible to have $C_{i(k)} = C_{i(k')}$ for $k \neq k'$).

Suppose first that $i(k) > j(k)$ for $k = 1, \ldots, \alpha$. Then, the subcurve $X_{j(k)}$ of lemma 2.4 is the connected component of $X - P_k$ which contains $C_{j(k)}$.

Let us see that $\overline{D} = \bigsqcup_{k=1}^{\alpha} X_{j(k)}$. Since the inclusion $\overline{D} \subseteq \bigcup_{k=1}^{\alpha} X_{j(k)}$ is clear we have only to prove that $\bigsqcup_{k=1}^{\alpha} X_{j(k)}$ contains no component of D and that $X_{j(k)} \cap X_{j(k')} = \emptyset$ if $k \neq k'$. If $C_{i(k')} \subseteq X_{j(k)}$ for some k', then $C_{j(k')} \subseteq X_{j(k)}$ and $j(k') < i(k') < j(k) < i(k)$. Hence,

$$X - P_k - P_{k'} = (Z_k^1 \sqcup Z_k^2) \sqcup \overline{X_{j(k)}}$$

where we denote by Z_k^1 (resp. Z_k^2) the connected component of $X_{j(k)} - P_{k'}$ that contains $C_{i(k')}$ (resp. $C_{j(k')}$). Analogously, $C_{j(k)}$, $C_{i(k)}$ are contained in $\overline{X_{j(k')}}$ and

$$X - P_{k'} - P_k = X_{j(k')} \sqcup (Z_{k'}^1 \sqcup Z_{k'}^2)$$

where $Z_{k'}^1$ (resp. $Z_{k'}^2$) is the connected component of $\overline{X_{j(k')}} - P_k$ containing $C_{i(k)}$ (resp. $C_{j(k)}$). We deduce that $X_{j(k')} = Z_k^2$, $\overline{X_{j(k)}} = Z_{k'}^1$ and $Z_{k'}^2 = Z_k^1$. Thus, $C_{j(k)} \subseteq Z_k^1$ and $j(k) < i(k')$, which is absurd. Therefore, none of the components $C_{i(k')}$ of D is in $\bigcup_{k=1}^{\alpha} X_{j(k)}$ and, since $X_{j(k)}$ are connected, no other component of D is either. Moreover, a similar argument shows that $X_{j(k)}$ and $X_{j(k')}$ have no common components for $k \neq k'$.

Let now $t_k + 1$ be the number of irreducible components in $X_{j(k)}$. Then, $\sum_k (t_k + 1)$ is equal to $N - t$. On the other hand, the number of intersection points in $\bigsqcup_k X_{j(k)}$ is $\sum_k t_k + s$, where s is the number of intersection points of $X_{j(k)}$, $k = 1, \ldots, \alpha$. Since $\sum_k t_k + s$ is the number $N - t - \alpha$ of intersection points in $\overline{D}$, we get $s = 0$ and $\overline{D} = \bigsqcup_{k=1}^{\alpha} X_{j(k)}$. Then, $d_{\overline{D}} = \sum_k d_{X_{j(k)}}$, $h_{\overline{D}} = \sum_k h_{X_{j(k)}}$ and $\chi(\mathcal{O}_{\overline{D}}) = \sum_k \chi(\mathcal{O}_{X_{j(k)}})$.

By definition of d_i^X, for every $k = 1, \ldots, \alpha$, we have

$$-\chi(\mathcal{O}_{X_{j(k)}}) + h_{X_{j(k)}} t + k_{X_{j(k)}} < d_{X_{j(k)}} < -\chi(\mathcal{O}_{X_{j(k)}}) + h_{X_{j(k)}} t + k_{X_{j(k)}} + 1 \tag{5}$$

Taking into account that $d = d_D + d_{\overline{D}}$, $h = h_D + h_{\overline{D}}$ and $\chi(\mathcal{O}_X) = \chi(\mathcal{O}_D) + \chi(\mathcal{O}_{\overline{D}}) - \alpha$, we obtain the result in this case.

Assume now that $j(1) > i(1)$. Then, $X_{i(1)}$ is the connected component of $X - P_1$ which contains $C_{i(1)}$. Since D is connected, $X_{i(1)}$ contains D and then $X_{i(1)}$ contains $C_{i(k)}$ and $C_{j(k)}$ for $k = 2, \ldots, \alpha$. This implies that $i(k) < i(1)$ for $k = 2, \ldots, \alpha$. If we had $i(k) < j(k)$ for some $k = 2, \ldots, \alpha$, then D would be contained in $X_{i(k)}$ as before and $i(1) < i(k)$, which is absurd. Thus, $i(k) > j(k)$ for $k = 2, \ldots, \alpha$. In this situation, one has that $X_{i(1)} = D \cup Z$, where $Z = \bigcup_{k=2}^{\alpha} X_{j(k)}$ and D intersects Z in $\alpha - 1$ points.

Thus, we have relations (5) for $k = 2, \ldots, \alpha$ and, arguing as in the former case, we obtain

$$-\chi(\mathcal{O}_Z) + h_Z t + k_Z < d_Z < -\chi(\mathcal{O}_Z) + h_Z t + k_Z + (\alpha - 1) \tag{6}$$

By definition of d_i^X, one has

$$-\chi(\mathcal{O}_{X_{i(1)}}) + h_{X_{i(1)}} t + k_{X_{i(1)}} < d_{X_{i(1)}} < -\chi(\mathcal{O}_{X_{i(1)}}) + h_{X_{i(1)}} t + k_{X_{i(1)}} + 1$$

which together with (6) proves the statement.

If the subcurve D is not connected, the inequality holds for every connected component and then it is easy to deduce it for D.

2. By lemma 2.2, it is enough to show that $\mu_H(L^D) < d$ for every proper subcurve D of X. Since

$$\mu_H(L^D) = \frac{hd - hd_D + h_D \chi(\mathcal{O}_X) - h\chi(\mathcal{O}_D)}{h - h_D},$$

the result follows from 1. □

Proof of Theorem 3.1. By lemmas 3.1 and 3.2, $\mathrm{Jac}^d(X)_s$ is not empty only if k_{X_i} is not integer for every $i \leq N - 1$, and in this case it is equal to $\prod_{i=1}^{N} \mathrm{Pic}^{d_i^X}(C_i)$.

We prove now the remaining statements of the theorem. If L is a strictly semistable line bundle on X of degree d then, hd_{X_i} is equal to one of the two extremal values of the inequality (4). In particular, k_{X_i} is an integer for some $i \leq N-1$.

Let i be the first index such that k_{X_i} is integer. Then, there are two possibilities for d_{X_i}:

a) $d_{X_i} = -\chi(\mathcal{O}_{X_i}) + h_{X_i}t + k_{X_i}$

b) $d_{X_i} = -\chi(\mathcal{O}_{X_i}) + h_{X_i}t + k_{X_i} + 1$

Let us construct a Jordan-Hölder filtration for L in both cases. Since case a) and case b) are the same but with the roles of X_i and $\overline{X_i}$ intertwined, we give the construction in the case a).

We have that $\mu_H(L_{X_i}) = \mu_H(L^{X_i}) = \mu_H(L)$. Then, L_{X_i} and $L^{X_i} \simeq L_{\overline{X_i}}(-P_i)$ are semistable with respect to H and, by lemma 2.1, they are semistable with respect to H_{X_i} and $H_{\overline{X_i}}$ respectively.

For simplicity, we shall write $Y = X_i = C_{i_0} \cup C_{i_1} \cup \ldots \cup C_{i_k}$ with $i_1, \ldots, i_k < i_0 = i$ and $Z = \overline{X_i}$, which are again generalized tree-like curves.

Let us see when the sheaves L_Y and $L_Z(-P_i)$ are stable. We can fix an ordering for Y as in lemma 2.4, so that $Y = C_{\sigma(i_0)} \cup \ldots \cup C_{\sigma(i_k)}$ and we obtain subcurves Y_r of Y for $r = \sigma(i_0), \ldots, \sigma(i_{k-1})$.

Claim 1. The sheaf L_Y is stable if and only if k_{Y_r} is not an integer for $r = \sigma(i_0), \ldots, \sigma(i_{k-1})$.

Proof. Since the residue class of $d_Y - \mathrm{g}(Y)$ modulo h_Y is $b_Y = k_Y - 1$, the numbers $\frac{h_{Y_r}(b_Y+1)}{h_Y} = k_{Y_r}$ are not integers for $r = \sigma(i_0), \ldots, \sigma(i_{k-1})$. Then, from lemma 3.2, we have only to prove that L_Y is in $\prod_r \operatorname{Pic}^{d_r^Y}(C_r)$, where d_r^Y are the integer numbers defined as d_i^X but with the new ordering of Y and r runs through the irreducible components of Y. This is equivalent to proving that

$$d_{Y_r} = -\chi(\mathcal{O}_{Y_r}) + h_{Y_r}t + [k_{Y_r}] + 1 \qquad \text{for } r = \sigma(i_0), \ldots, \sigma(i_{k-1}).$$

Actually, since L is semistable and Y_r is a proper subcurve of X, arguing as in lemma 3.2, we obtain

$$-\chi(\mathcal{O}_{Y_r}) + h_{Y_r}t + k_{Y_r} \leq d_{Y_r} \leq -\chi(\mathcal{O}_{Y_r}) + h_{Y_r}t + k_{Y_r} + \alpha \tag{7}$$

where α is the number of intersection points of Y_r and its complement in X. We have that $\alpha \leq 2$ and d_{Y_r} is not equal to the extremal values of (7) because $k_{Y_r} \notin \mathbb{Z}$. Moreover, if it were

$$d_{Y_r} = -\chi(\mathcal{O}_{Y_r}) + h_{Y_r}t + [k_{Y_r}] + 2,$$

since $d_Y = d_{Y_r} + d_{\overline{Y_r}^Y}$, $h_Y = h_{Y_r} + h_{\overline{Y_r}^Y}$ and $\chi(\mathcal{O}_Y) = \chi(\mathcal{O}_{Y_r}) + \chi(\mathcal{O}_{\overline{Y_r}^Y}) - 1$, $\overline{Y_r}^Y$ being the complement of Y_r in Y, then

$$d_{\overline{Y_r}^Y} = -\chi(\mathcal{O}_{\overline{Y_r}^Y}) + h_{\overline{Y_r}^Y} t + [k_{\overline{Y_r}^Y}]$$

which contradicts the semistability of L. Thus,

$$d_{Y_r} = -\chi(\mathcal{O}_{Y_r}) + h_{Y_r} t + [k_{Y_r}] + 1$$

and the proof of the claim 1 is complete.

On the other hand, the irreducible components of Z are ordered according the instructions in lemma 2.4 and the subcurves Z_s, where s runs through the irreducible components of Z and $s \leq N-1$, are equal to either X_s or $X_s - Y$.

Claim 2. The sheaf $L_Z(-P_i)$ is stable if and only if k_{X_s} is not an integer for every $s > i$.

Proof. Since the residue class of $d_Z - 1 - \mathrm{g}(Z)$ modulo h_Z is $b_Z = k_Z - 1$ and $k_Y \in \mathbb{Z}$, by the hypothesis, the numbers $\frac{h_{Z_s}(b_Z+1)}{h_Z}$ are not integers for $s > i$ and, by the choice of i, they aren't for $s < i$ either. Then, by lemma 3.2, it is enough to prove that

$$d_{H_{Z_s}}(L_Z(-P_i)|_{Z_s}) = -\chi(\mathcal{O}_{Z_s}) + h_{Z_s} t + [k_{Z_s}] + 1 \quad \text{for } s \leq N-1.$$

Since L is semistable, we have that

$$d_{X_s} = -\chi(\mathcal{O}_{X_s}) + h_{X_s} t + [k_{X_s}] + 1.$$

Moreover, if $Z_s = X_s$ then, $d_{H_{Z_s}}(L_Z(-P_i)|_{Z_s}) = d_{X_s}$ and if $Z_s = X_s - Y$ then, $d_{H_{Z_s}}(L_Z(-P_i)|Z_s) = d_{X_s} - d_Y - 1$. We obtain the desired result in both cases and the proof of the claim 2 is complete.

We return now to the proof of the theorem. If k_{Y_r} and k_{X_s} are not integers for $r = \sigma(i_0), \ldots, \sigma(i_{k-1})$ and $s > i$, then $0 \subset L_Z(-P_i) \subset L$ is a Jordan-Hölder filtration for L and the S-equivalence class of L belongs to $\prod_r \mathrm{Pic}^{d_r^Y}(C_r) \times \prod_s \mathrm{Pic}^{d_s^X}(C_s)$.

On the other hand, if k_{Y_r} is integer for some $r = \sigma(i_0), \ldots, \sigma(i_{k-1})$, the sheaf L_Y is strictly semistable and we have to repeat the above procedure with L_Y in the place of L and the curve Y in the place of X. Similarly, if k_{X_s} is integer for some $s > i$, the sheaf $L_Z(-P_i)$ is strictly semistable. Then, we have to repeat the above procedure for $L_Z(-P_i)$. By iterating this procedure, we get a Jordan-Hölder filtration for L_Y:

$$0 = F_0 \subset F_1 \subset \ldots \subset F_m = L_Y$$

and another for $L_Z(-P_i)$:

$$0 = G_0 \subset G_1 \subset \ldots \subset G_n = L_Z(-P_i).$$

Therefore, a filtration for L is given by

$$0 = G_0 \subset G_1 \subset \ldots \subset L_Z(-P_i) \subset \pi_Y^{-1}(F_1) \subset \ldots \subset \pi_Y^{-1}(L_Y) = L.$$

Thus, the S-equivalence class of L belongs to $\prod_{i=1}^N \operatorname{Pic}^{d_i}(C_i)$, where d_i are the integer numbers constructed with the algorithm.

Finally, let us consider a torsion free sheaf F on X of rank 1 and degree d which is not locally free. When F is locally free at the intersection points P_i for all $i = 1, \ldots, N-1$, calculations and results are analogous to the former ones. If F is not locally free at P_i for some $i = 1, \ldots, N-1$, then there is a natural morphism

$$F \to F_Y \oplus F_Z$$

where Y, Z are the connected components of $X - P_i$, that is clearly an isomorphism outside P_i. But this is an isomorphism at P_i as well because by lemma 2.3, $F_{P_i} \simeq \mathcal{O}_{C_i^1, P_i} \oplus \mathcal{O}_{C_i^2, P_i}$ and this is precisely the stalk of $F_Y \oplus F_Z$ at P_i. We conclude that if F is strictly semistable, then k_Y y k_Z are integers, d_Y and d_Z are given by

$$d_Y = -\chi(\mathcal{O}_Y) + h_Y t + k_Y, \qquad d_Z = -\chi(\mathcal{O}_Z) + h_Z t + k_Z,$$

and F_Y and F_Z are semistable as well. Then, the construction of a Jordan-Hölder filtration for F can be done as above and thus the S-equivalence class of F belongs to $\prod_{i=1}^N \overline{\operatorname{Jac}}^{d_i}(C_i)$. $\square$

We now give three examples to illustrate this theorem.

Example 1. Let $X = C_1 \cup \ldots \cup C_N$, $N \geq 2$, be a generalized tree-like curve with a polarization H whose degree h is a prime number. Suppose that the irreducible components of X are ordered according with lemma 2.4. Then, since h_{X_i} is not divisible by h, $k_{X_i} = \frac{h_{X_i}(b+1)}{h}$ is an integer if and only if $b = h - 1$. Therefore, by theorem 3.1, we have that if $b < h - 1$, $\mathrm{Jac}^d(X)_s = \prod_{i=1}^N \mathrm{Pic}^{d_i^X}(C_i)$ and

$$\mathrm{Jac}^d(X)_s \subsetneq \overline{\mathrm{Jac}}^d(X)_s = \overline{\mathrm{Jac}}^d(X) \simeq \prod_{i=1}^N \overline{\mathrm{Jac}}^{d_i^X}(C_i),$$

whereas for $b = h - 1$, $\mathrm{Jac}^d(X)_s$ and $\overline{\mathrm{Jac}}^d(X)_s$ are empty and

$$\overline{\mathrm{Jac}}^d(X) \simeq \prod_{i=1}^N \overline{\mathrm{Jac}}^{d_i}(C_i).$$

Moreover, in this case the number d_i is given by

$$d_i = h_{C_i}(t+1) - \chi(\mathcal{O}_{C_i}) \quad \text{for } i = 1, \ldots, N.$$

Actually, if L is a strictly semistable line bundle of degree d on X, the first index i such that k_{X_i} is integer is $i = 1$ and the connected components of $X - P_1$ are $Y = C_1$ and $Z = C_2 \cup \ldots \cup C_N$. Suppose, as in the proof of the theorem, that

$$d_Y = -\chi(\mathcal{O}_Y) + h_Y t + k_Y = h_Y(t+1) - \chi(\mathcal{O}_Y).$$

Then, L_Y and $L_Z(-P_1)$ are semistable and $L_Y = L_{C_1} \in \mathrm{Pic}^{h_{C_1}(t+1)-\chi(\mathcal{O}_{C_1})}(C_1)$. On the other hand, since k_{X_s} is integer for $s = 2, \ldots, N$, we have to apply the procedure to the curve Z. Here, the first index s such that k_{X_s} is integer is $s = 2$ and the connected components of $Z - P_2$ are C_2 and $C_3 \cup \ldots \cup C_N$. Proceeding exactly as in case of X, we obtain a sheaf belonging to $\mathrm{Pic}^{h_{C_2}(t+1)-\chi(\mathcal{O}_{C_2})}(C_2)$ and another supported on $C_3 \cup \ldots \cup C_N$. Since k_{X_s} is integer for $s = 3, \ldots, N$, we apply the procedure to the curve $C_3 \cup \ldots \cup C_N$, and so on. The iteration of this procedure will only finish when we obtain a sheaf supported on C_N that belongs to $\mathrm{Pic}^{h_{C_N}(t+1)-\chi(\mathcal{O}_{C_N})}(C_N)$. Thus, $d_i = h_{C_i}(t+1) - \chi(\mathcal{O}_{C_i})$ for $i = 1, \ldots, N$.

Example 2. We are now going to recover examples 1 and 2 of [A]. There, the irreducible components C_i are taken to be smooth and $d = g - 1$. Then, the residue class of $d - g$ modulo h is $b = h - 1$ and $t = -1$. It follows that k_{X_i} is integer for $i = 1, \ldots, N$ and, arguing as in our example 1, we have that $\mathrm{Jac}^{g-1}(X)_s$ and $\overline{\mathrm{Jac}}^{g-1}(X)_s$ are empty and

$$\overline{\mathrm{Jac}}^{g-1}(X) \simeq \prod_{i=1}^N \overline{\mathrm{Jac}}^{h_{C_i}(t+1)-\chi(\mathcal{O}_{C_i})}(C_i) \simeq \prod_{i=1}^N \mathrm{Pic}^{g_i-1}(C_i)$$

as asserted in [A].

Example 3. Let X be the following tree-like curve



Fixing an ordering of the irreducible components of X as in lemma 2.4, we obtain



so that $X_1 = C_1$, $X_2 = C_2$, $X_3 = C_1 \cup C_2 \cup C_3$. Assume that the first index i such that k_{X_i} is integer is $i = 3$ and let us compute the numbers d_i, $i = 1, \ldots, 4$ in this case.

The connected components of $X - P_3$ are $Y = C_1 \cup C_2 \cup C_3$ and $Z = C_4$. Suppose L is a strictly semistable line bundle of degree d on X and that

$$d_Y = -\chi(\mathcal{O}_Y) + h_Y t + k_Y$$

(the other case is similar). Then L_Y and $L_Z(-P_3)$ are semistable. Moreover, $L_Z(-P_3) \in \mathrm{Pic}^{h_{C_4} t + k_{C_4} - \chi(\mathcal{O}_{C_4})}(C_4)$ so that

$$d_4 = h_{C_4} t + k_{C_4} - \chi(\mathcal{O}_{C_4}).$$

We now have to fix a new ordering for Y according with lemma 2.4. We can take, for instance, $Y = C_{\sigma(1)} \cup C_{\sigma(2)} \cup C_{\sigma(3)}$ with $\sigma(1) = 1$, $\sigma(2) = 3$ and $\sigma(3) = 2$. Then, $Y_{\sigma(1)} = C_1$ and $Y_{\sigma(2)} = C_1 \cup C_3$. Therefore, since $k_{Y_{\sigma(1)}} = k_{X_1}$

and $k_{Y_{\sigma(2)}} = k_Y - k_{X_2}$ are not integers, we conclude that

$$\begin{aligned}
&d_1 = d^Y_{\sigma(1)} = -\chi(\mathcal{O}_{Y_{\sigma(1)}}) + h_{Y_{\sigma(1)}} t + [k_{Y_{\sigma(1)}}] + 1 = d^X_1,\\
&d_3 = d^Y_{\sigma(2)} = -\chi(\mathcal{O}_{Y_{\sigma(2)}}) + h_{Y_{\sigma(2)}} t + [k_{Y_{\sigma(2)}}] + 1 - d^Y_{\sigma(1)} =\\
&\quad = -\chi(\mathcal{O}_Y) + h_Y t + k_Y + 1 - d^X_1 - d^X_2 - 1 = d^X_3 - 1,\\
&d_2 = d^Y_{\sigma(3)} = d_Y - d^Y_{\sigma(1)} - d^Y_{\sigma(2)} = d^X_2.
\end{aligned}$$

References

[A] V. Alexeev, *Compactified Jacobians* Preprint alg-geom/9608012 (1996).

[AK] A. Altman, S. Kleiman, *Compactifying the Picard Scheme*, Adv. in Math., **35** (1980), 50-112.

[B] A. Beauville, *Counting rational curves on K3 surfaces*, Preprint alg-geo/9701019 (1997).

[C] L. Caporaso, *A compactification of the universal Picard variety over the moduli space of stable curves*, J. Amer. Math. Soc **7** (1994), 589-660.

[D'So] C. D'Souza, *Compactification of generalized Jacobians*, Proc. Indian Acad. Sci. Sect. A Math. Sci. **88** (1979), 419-457.

[E] E. Esteves, *Compactifying the relative Jacobian over families of reduced curves*, Preprint alg-geom/9709009 (1997).

[HM] D. Hernández Ruipérez, J.M. Muñoz Porras *Structure of the moduli space of stable sheaves on elliptic fibrations*, Preprint Univ. de Salamanca (1999).

[I] J. Igusa, *Fiber systems of Jacobian varieties*, Amer. J. Math., **78** (1956), 171-199.

[OS] T. Oda, C. S. Seshadri, *Compactifications of the generalized Jacobian variety*, Trans. Amer. Math. Soc., **253** (1979), 1-90.

[P] R. Pandharipande, *A compactification over* $\overline{M}_g$ *of the universal moduli space of slope-semistable vector bundles*, J. Amer. Math. Soc., **9** (1996), 425-471.

[S] C. T. Simpson, *Moduli of Representations of Fundamental Group of a Smooth Projective Variety I*, Publ. Math. Inst. Hautes Études Sci. **79** (1994), 47-129.

[Ses] C.S. Seshadri, *Fibrés vectoriels sur les courbes algébriques*, Astérisque **96** (1982).

[T] M. Teixidor, *Moduli spaces of (semi)stable vector bundles on tree-like curves*, Math. Ann. **290** (1991), 341-348.